

SOME INTEGRAL INEQUALITIES FOR THE POLAR DERIVATIVE OF A POLYNOMIAL

Abdullah Mir

(Kashmir University, India)

Sajad Amin Baba

(Govt. Hr. Sec. Institute, India)

Received Oct. 9, 2010

© Editorial Board of Analysis in Theory & Applications and Springer-Verlag Berlin Heidelberg 2011

Abstract. If $P(z)$ is a polynomial of degree n which does not vanish in $|z| < 1$, then it is recently proved by Rather [*Jour. Ineq. Pure and Appl. Math.*, 9 (2008), Issue 4, Art. 103] that for every $\gamma > 0$ and every real or complex number α with $|\alpha| \geq 1$,

$$\left\{ \int_0^{2\pi} |D_\alpha P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma} \leq n(|\alpha| + 1)C_\gamma \left\{ \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma},$$

$$C_\gamma = \left\{ \frac{1}{2\pi} \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta \right\}^{-1/\gamma},$$

where $D_\alpha P(z)$ denotes the polar derivative of $P(z)$ with respect to α . In this paper we prove a result which not only provides a refinement of the above inequality but also gives a result of Aziz and Dawood [*J. Approx. Theory*, 54 (1988), 306-313] as a special case.

Key words: polar derivative, polynomial, Zygmund inequality, zeros

AMS (2010) subject classification: 30A10, 30C10, 30D15, 41A17

1 Introduction and Statement of Results

Let $P(z) = \sum_{\nu=0}^n a_\nu z^\nu$ be a polynomial of degree at most n and $P'(z)$ its derivative, then

$$\max_{|z|=1} |P'(z)| \leq n \max_{|z|=1} |P(z)|, \tag{1.1}$$

and for every $\gamma \geq 1$,

$$\left\{ \int_0^{2\pi} |P'(e^{i\theta})|^\gamma r m d\theta \right\}^{1/\gamma} \leq n \left\{ \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma}. \tag{1.2}$$

The inequality (1.1) is a classical result of Bernstein^[11] (see also [14]), whereas the inequality (1.2) is due to Zygmund^[15], who proved it for all trigonometric polynomials of degree n and not only for those of the form $P(e^{i\theta})$. Arestov^[1] proved that (1.2) remains true for $0 < \gamma < 1$ as well. If we let $\gamma \rightarrow \infty$ in the inequality (1.2), we get (1.1).

The above two inequalities (1.1) and (1.2) can be sharpened if we restrict ourselves to the class of polynomials having no zeros in $|z| < 1$. In fact, if $P(z) \neq 0$ in $|z| < 1$, then (1.1) and (1.2) can be respectively replaced by

$$\max_{|z|=1} |P'(z)| \leq \frac{n}{2} \max_{|z|=1} |P(z)| \tag{1.3}$$

and

$$\left\{ \int_0^{2\pi} |P'(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma} \leq n B_\gamma \left\{ \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma}, \tag{1.4}$$

where

$$B_\gamma = \left\{ \frac{1}{2\pi} \int_0^{2\pi} |1 + e^{i\alpha}|^\gamma d\alpha \right\}^{-1/\gamma}.$$

The inequality (1.3) is conjectured by Erdős and later verified by Lax^[9], whereas the inequality (1.4) is proved by De-Bruijn^[7] for $\gamma \geq 1$. Further, Rahman and Schmeisser^[12] have shown that (1.4) holds for $0 < \gamma < 1$ also. If we let $\gamma \rightarrow \infty$ in the inequality (1.4), we get (1.3).

The inequality (1.3) is further improved by Aziz and Dawood^[4] by proving that if $P(z) \neq 0$ in $|z| < 1$, then

$$\max_{|z|=1} |P'(z)| \leq \frac{n}{2} \left\{ \max_{|z|=1} |P(z)| - \min_{|z|=1} |P(z)| \right\}. \tag{1.5}$$

Let $D_\alpha P(z)$ denote the polar derivative of the polynomial $P(z)$ with respect to a complex number α . Then

$$D_\alpha P(z) = nP(z) + (\alpha - z)P'(z).$$

The polynomial $D_\alpha P(z)$ is of degree at most $n - 1$ and it generalizes the ordinary derivative $P'(z)$ in the sense that

$$\lim_{\alpha \rightarrow \infty} \frac{D_\alpha P(z)}{\alpha} = P'(z).$$

Aziz^[3] extended the inequality (1.3) to the polar derivatives and proved that if $P(z)$ is a polynomial of degree n such that $P(z) \neq 0$ in $|z| < 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha P(z)| \leq \frac{n}{2} (|\alpha| + 1) \max_{|z|=1} |P(z)|. \tag{1.6}$$

While seeking the desired extension of the inequality (1.6) to the L^γ norm, recently Govil et al. [8] have made an incomplete attempt by proving the following generalization of the inequalities (1.4) and (1.6).

Theorem A. *If $P(z)$ is a polynomial of degree n which does not vanish in $|z| < 1$, then for $\gamma \geq 1$ and every real or complex number α with $|\alpha| \geq 1$,*

$$\left\{ \int_0^{2\pi} |D_\alpha P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma} \leq n(|\alpha| + 1)F_\gamma \left\{ \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma}, \tag{1.7}$$

where

$$F_\gamma = \left\{ \frac{1}{2\pi} \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta \right\}^{-1/\gamma}.$$

Unfortunately, the proof of Theorem A is not correct as is first pointed out by Aziz and Rather^[5] who in the same paper have given a correct proof of the inequality (1.7) also. The inequality (1.7) is then independently proved by Rather^[13] for $\gamma > 0$.

In this paper we prove the following more general result which in particular provides refinements and generalizations of the inequalities (1.6) and (1.7) and also extends the inequality (1.7) for $\gamma \in (0, 1)$. Further, it also gives the inequality (1.5) as a special case. Actually, we prove

Theorem 1.1. *If $P(z)$ is a polynomial of degree n which does not vanish in $|z| < 1$, then for $\gamma > 0$, every real or complex numbers $\alpha_1, \dots, \alpha_k$, $k \leq n - 1$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$ and real or complex δ with $|\delta| \leq 1$,*

$$\left\{ \int_0^{2\pi} \left| D_{\alpha_1} \dots D_{\alpha_k} P(e^{i\theta}) + \frac{mn(n-1) \dots (n-k+1)(|\alpha_1 \dots \alpha_k| - 1)\delta}{2} \right|^\gamma d\theta \right\}^{1/\gamma} \leq n(n-1) \dots (n-k+1)(|\alpha_1| + 1)(|\alpha_2| + 1) \dots (|\alpha_k| + 1)C_\gamma \left\{ \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta \right\}^{1/\gamma}, \tag{1.8}$$

where

$$C_\gamma = \left\{ \frac{1}{2\pi} \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta \right\}^{-1/\gamma}$$

and

$$m = \min_{|z|=1} |P(z)|.$$

In the limiting case, when $\gamma \rightarrow \infty$, the above inequality is sharp and the equality in (1.8) holds for $P(z) = (z + 1)^n$, where $\alpha_i \geq 1$, $i = 1, 2, \dots, k$ are real.

If we let $\gamma \rightarrow \infty$ in (1.8) and choose the argument of δ with $|\delta| = 1$ suitably, we get the following refinement and generalization of (1.6).

Corollary 1.1. *If $P(z)$ is a polynomial of degree n such that $P(z) \neq 0$ in $|z| < 1$, then for every real or complex numbers $\alpha_1, \dots, \alpha_k$, $k \leq n - 1$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$*

$$\max_{|z|=1} |D_{\alpha_1} \cdots D_{\alpha_k} P(z)| \leq \frac{n(n-1) \cdots (n-k+1)}{2} \left\{ (|\alpha_1| + 1) \cdots (|\alpha_k| + 1) \max_{|z|=1} |P(z)| - (|\alpha_1 \cdots \alpha_k| - 1) \min_{|z|=1} |P(z)| \right\}. \tag{1.9}$$

The result is best possible and the equality holds in (1.9) for $P(z) = (z + 1)^n$ with real $\alpha_i \geq 1$, $i = 1, 2, \dots, k$.

If we put $k = 1$, in Theorem 1.1, we get the following result which is a refinement of (1.7) and is an extension for $\gamma \in (0, 1)$.

Corollary 1.2. *If $P(z)$ is a polynomial of degree n which does not vanish in $|z| < 1$, then for $\gamma > 0$, every real or complex number α with $|\alpha| \geq 1$ and real or complex δ with $|\delta| \leq 1$,*

$$\left\{ \int_0^{2\pi} \left| D_{\alpha} P(e^{i\theta}) + \frac{m n (|\alpha| - 1) \delta}{2} \right|^{\gamma} d\theta \right\}^{1/\gamma} \leq n (|\alpha| + 1) C_{\gamma} \left\{ \int_0^{2\pi} |P(e^{i\theta})|^{\gamma} d\theta \right\}^{1/\gamma}, \tag{1.10}$$

where C_{γ} , m are defined above. In the limiting case, when $\gamma \rightarrow \infty$, the above inequality is sharp and the equality in (1.10) holds for $P(z) = (z + 1)^n$, where $\alpha \geq 1$ is real.

If we let $\gamma \rightarrow \infty$ in (1.10) and choose the argument of δ with $|\delta| = 1$ suitably, we get the following refinement of (1.6).

Corollary 1.3. *If $P(z)$ is a polynomial of degree n such that $P(z) \neq 0$ in $|z| < 1$, then for every real or complex number α with $|\alpha| \geq 1$,*

$$\max_{|z|=1} |D_{\alpha} P(z)| \leq \frac{n}{2} \left\{ (|\alpha| + 1) \max_{|z|=1} |P(z)| - (|\alpha| - 1) \min_{|z|=1} |P(z)| \right\}. \tag{1.11}$$

The result is best possible and the equality holds in (1.11) for $P(z) = (z + 1)^n$ with real $\alpha \geq 1$.

Remark 1.1. If we divide both sides of (1.11) by $|\alpha|$ and let $|\alpha| \rightarrow \infty$, we get (1.5).

2 Lemmas

We need the following lemmas for the proof of Theorem 1.1.

Lemma 2.1. *If all the zeros of an n th degree polynomial $P(z)$ lie in a circular region C and if none of the points $\alpha_1, \alpha_2, \dots, \alpha_k$ lie in the region C , then each of the polar derivatives*

$$D_{\alpha_1} \cdots D_{\alpha_k} P(z), \quad k = 1, 2, \dots, n - 1,$$

has all of its zeros in C .

This follows by repeated application of Laguarre’s theorem (see [1] or [9, p.52]).

Lemma 2.2. *If $P(z)$ is a polynomial of degree n having no zeros in $|z| < 1$ and $m = \min_{|z|=1} |P(z)|$, then for any real or complex numbers $\alpha_1, \dots, \alpha_k$, $k \leq n - 1$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$*

$$|D_{\alpha_1} \cdots D_{\alpha_k} Q(z)| \geq mn(n - 1) \cdots (n - k + 1) |\alpha_1 \alpha_2 \cdots \alpha_k z^{n-k}|, \quad \text{for } |z| \geq 1, \tag{2.1}$$

where

$$Q(z) = z^n \overline{P(1/\bar{z})}.$$

Proof of Lemma 2.2. If $m = \min_{|z|=1} |P(z)| = 0$, then the inequality (2.1) is obvious. Henceforth, we assume $m \neq 0$, so that all zeros of $P(z)$ lie in $|z| > 1$. Now if λ is any real or complex number with $|\lambda| < 1$, then

$$|\lambda m| < m \leq |P(z)|, \quad \text{for } |z| = 1. \tag{2.2}$$

Therefore, it follows by Rouché’s theorem that the polynomial $F(z) = P(z) - \lambda m$ has all zeros in $|z| > 1$ for every λ with $|\lambda| < 1$.

If $G(z) = z^n \overline{F(1/\bar{z})} = Q(z) - \bar{\lambda} m z^n$, then all zeros of $G(z)$ lie in $|z| < 1$. Hence, it follows by Lemma 2.1 that all zeros of

$$\begin{aligned} &D_{\alpha_1} \cdots D_{\alpha_k} (Q(z) - \bar{\lambda} m z^n) \\ &= D_{\alpha_1} \cdots D_{\alpha_k} Q(z) - \bar{\lambda} mn(n - 1) \cdots (n - k + 1) \alpha_1 \alpha_2 \cdots \alpha_k z^{n-k} \end{aligned} \tag{2.3}$$

lie in $|z| < 1$ for any $\alpha_1, \dots, \alpha_k$, $k \leq n - 1$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$ and for every λ with $|\lambda| < 1$. This implies

$$|D_{\alpha_1} \cdots D_{\alpha_k} Q(z)| \geq mn(n - 1) \cdots (n - k + 1) |\alpha_1 \alpha_2 \cdots \alpha_k z^{n-k}|, \quad \text{for } |z| \geq 1,$$

because if this is not true, then there is a point $z = z_0$ with $|z_0| \geq 1$, such that

$$|D_{\alpha_1} \cdots D_{\alpha_k} Q(z)|_{z=z_0} < mn(n - 1) \cdots (n - k + 1) |\alpha_1 \alpha_2 \cdots \alpha_k z_0^{n-k}|.$$

We take

$$\bar{\lambda} = \frac{\{D_{\alpha_1} \cdots D_{\alpha_k} Q(z)\}_{z=z_0}}{mn(n - 1) \cdots (n - k + 1) \alpha_1 \alpha_2 \cdots \alpha_k z_0^{n-k}},$$

so that $|\lambda| < 1$ and from (2.3) with this choice of $\bar{\lambda}$, we get $[D_{\alpha_1} \cdots D_{\alpha_k} (Q(z) - \bar{\lambda} m z^n)]_{z=z_0} = 0$, where $|z_0| \geq 1$, which contradicts the fact that all zeros of $D_{\alpha_1} \cdots D_{\alpha_k} (Q(z) - \bar{\lambda} m z^n)$ lie in $|z| < 1$ and this completes the proof of lemma 2.2.

Lemma 2.3. *If $P(z)$ is a polynomial of degree n having no zeros in $|z| < 1$ and $m = \min_{|z|=1} |P(z)|$, $Q(z) = z^n \overline{P(1/\bar{z})}$, then for any real or complex numbers $\alpha_1, \dots, \alpha_k$, $k \leq n - 1$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$ we have*

$$\begin{aligned} & |D_{\alpha_1} \cdots D_{\alpha_k} P(z)| \\ & \leq |D_{\alpha_1} \cdots D_{\alpha_k} Q(z)| - mn(n-1) \cdots (n-k+1) (|\alpha_1 \alpha_2 \cdots \alpha_k| - 1), \text{ for } |z| = 1. \end{aligned}$$

Proof of Lemma 2.3. Since $P(z)$ has all zeros in $|z| \geq 1$ and $m = \min_{|z|=1} |P(z)|$, then

$$m \leq |P(z)|, \text{ for } |z| = 1.$$

Therefore, for every real or complex number λ with $|\lambda| < 1$, it follows by Rouché's theorem for $m > 0$ that the polynomial $F(z) = P(z) - \lambda m$ has all zeros in $|z| > 1$ and hence no zero in $|z| < 1$. Thus the polynomial $T(z) = z^n \overline{F(1/\bar{z})} = Q(z) - \bar{\lambda} m z^n$ has all zeros in $|z| < 1$ and

$$|F(z)| \leq |T(z)|, \text{ for } |z| = 1.$$

It follows again by Rouché's theorem that for every β , $|\beta| > 1$, the polynomial $F(z) - \beta T(z)$ has all zeros in $|z| < 1$ which implies by Lemma 2.1 that for every real or complex numbers $\alpha_1, \dots, \alpha_k$ with $|\alpha_i| \geq 1$, $i = 1, 2, \dots, k$ the polynomial $D_{\alpha_1} \cdots D_{\alpha_k} [F(z) - \beta T(z)]$ has all zeros in $|z| < 1$. This implies

$$|D_{\alpha_1} \cdots D_{\alpha_k} F(z)| \leq |D_{\alpha_1} \cdots D_{\alpha_k} T(z)|, \text{ for } |z| \geq 1, \tag{2.4}$$

The inequality (2.4) is clearly equivalent to

$$|D_{\alpha_1} \cdots D_{\alpha_k} (P(z) - \lambda m)| \leq |D_{\alpha_1} \cdots D_{\alpha_k} (Q(z) - \bar{\lambda} m z^n)|, \text{ for } |z| \geq 1.$$

Equivalently,

$$\begin{aligned} & |D_{\alpha_1} \cdots D_{\alpha_k} P(z) - \lambda mn(n-1) \cdots (n-k+1)| \\ & \leq |D_{\alpha_1} \cdots D_{\alpha_k} Q(z) - \bar{\lambda} mn(n-1) \cdots (n-k+1) \alpha_1 \alpha_2 \cdots \alpha_k z^{n-k}|, \end{aligned}$$

which gives

$$\begin{aligned} & |D_{\alpha_1} \cdots D_{\alpha_k} P(z) - mn(n-1) \cdots (n-k+1) \lambda| \\ & \leq |D_{\alpha_1} \cdots D_{\alpha_k} Q(z) - \bar{\lambda} mn(n-1) \cdots (n-k+1) \alpha_1 \alpha_2 \cdots \alpha_k z^{n-k}|, \end{aligned} \tag{2.5}$$

for $|z| \geq 1$ and for every λ with $|\lambda| < 1$.

Now choosing the argument of λ suitably, so that on $|z| = 1$,

$$\begin{aligned} & |D_{\alpha_1} \cdots D_{\alpha_k} Q(z) - \bar{\lambda} mn(n-1) \cdots (n-k+1) \alpha_1 \alpha_2 \cdots \alpha_k z^{n-k}| \\ & = |D_{\alpha_1} \cdots D_{\alpha_k} Q(z) - mn(n-1) \cdots (n-k+1) \alpha_1 \alpha_2 \cdots \alpha_k| |\lambda|, \end{aligned} \tag{2.6}$$

we get from (2.5) that on $|z| = 1$,

$$\begin{aligned}
 &|D_{\alpha_1} \cdots D_{\alpha_k} Q(z)| \\
 &\geq |D_{\alpha_1} \cdots D_{\alpha_k} P(z)| + |\lambda| mn(n-1) \cdots (n-k+1) (|\alpha_1 \alpha_2 \cdots \alpha_k| - 1).
 \end{aligned}
 \tag{2.7}$$

The fact that the right hand side of (2.6) is non-negative follows from Lemma 2.2. Lemma 2.3 now follows by making $|\lambda| \rightarrow 1$ in (2.7).

Lemma 2.4. *If $P(z)$ is a polynomial of degree n then for every complex number α and $\gamma > 0$,*

$$\begin{aligned}
 &\left\{ \int_0^{2\pi} |D_{\alpha} P(e^{i\theta})|^{\gamma} d\theta \right\}^{1/\gamma} \\
 &\leq n(|\alpha| + 1) \left\{ \int_0^{2\pi} |P(e^{i\theta})|^{\gamma} d\theta \right\}^{1/\gamma}.
 \end{aligned}$$

Lemma 2.5. *If $P(z)$ is a polynomial of degree n which does not vanish in $|z| < t, t \geq 1$ and $Q(z) = z^n \overline{P(1/\bar{z})}$, then for every real or complex number α , real β with $0 \leq \beta < 2\pi$ and $\gamma > 0$,*

$$\begin{aligned}
 &\int_0^{2\pi} \int_0^{2\pi} |D_{\alpha} P(e^{i\theta}) + e^{i\beta} t^2 D_{\alpha/t^2} Q(e^{i\theta})|^{\gamma} d\theta d\beta \\
 &\leq 2\pi n^{\gamma} (|\alpha| + t)^{\gamma} \int_0^{2\pi} |P(e^{i\theta})|^{\gamma} d\theta.
 \end{aligned}$$

The above two lemmas are due to Rather^[12].

Lemma 2.6. *If A, B and C are non-negative real numbers such that $B + C \leq A$, then for every real number α ,*

$$|(A - C)e^{i\alpha} + (B + C)| \leq |Ae^{i\alpha} + B|.$$

This lemma is due to Aziz and Rather^[5].

3 Proof of Theorems

Proof of Theorem 1.1. Since $P(z)$ is a polynomial of degree at most n and $Q(z) = z^n \overline{P(1/\bar{z})}$, therefore for each $\beta, 0 \leq \beta < 2\pi, F(z) = P(z) + e^{i\beta} Q(z)$ is a polynomial of degree at most n so that $D_{\alpha_1} \cdots D_{\alpha_k} F(z) = D_{\alpha_1} \cdots D_{\alpha_k} P(z) + e^{i\beta} D_{\alpha_1} \cdots D_{\alpha_k} Q(z)$ is a polynomial of degree at most $n - k, k = 1, 2, \dots, n - 1$. By repeated application of Lemma 2.4, we have for each $\gamma > 0$,

$$\int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_k} F(e^{i\theta}) \right|^{\gamma} d\theta \leq (n - k + 1)^{\gamma} (|\alpha_k| + 1)^{\gamma} \int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_{k-1}} F(e^{i\theta}) \right|^{\gamma} d\theta.
 \tag{3.1}$$

Equivalently,

$$\begin{aligned}
 & \int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta}) \right|^\gamma d\theta \\
 & \leq (n-k+1)^\gamma (|\alpha_k|+1)^\gamma \int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_{k-1}} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} \cdots D_{\alpha_{k-1}} Q(e^{i\theta}) \right|^\gamma d\theta \\
 & \leq (n-k+1)^\gamma (n-k+2)^\gamma (|\alpha_k|+1)^\gamma (|\alpha_{k-1}|+1)^\gamma \\
 & \quad \times \int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_{k-2}} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} \cdots D_{\alpha_{k-2}} Q(e^{i\theta}) \right|^\gamma d\theta \tag{3.2} \\
 & \quad \vdots \\
 & \quad \vdots \\
 & \leq (n-k+1)^\gamma \cdots (n-1)^\gamma (|\alpha_k|+1)^\gamma \cdots (|\alpha_2|+1)^\gamma \\
 & \quad \times \int_0^{2\pi} \left| D_{\alpha_1} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} Q(e^{i\theta}) \right|^\gamma d\theta. \tag{3.3}
 \end{aligned}$$

Integrating both sides of (3.1) with respect to β from 0 to 2π , we get with the help of Lemma 2.5 (for $t = 1$) that for each $\gamma > 0$,

$$\begin{aligned}
 & \int_0^{2\pi} \int_0^{2\pi} \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta}) \right|^\gamma d\theta d\beta \\
 & \leq (n-k+1)^\gamma \cdots (n-1)^\gamma (|\alpha_k|+1)^\gamma \cdots (|\alpha_2|+1)^\gamma \int_0^{2\pi} \int_0^{2\pi} \left| D_{\alpha_1} P(e^{i\theta}) + e^{i\beta} D_{\alpha_1} Q(e^{i\theta}) \right|^\gamma d\theta d\beta \\
 & \leq 2\pi n^\gamma (n-1)^\gamma \cdots (n-k+1)^\gamma (|\alpha_k|+1)^\gamma \cdots (|\alpha_1|+1)^\gamma \int_0^{2\pi} \left| P(e^{i\theta}) \right|^\gamma d\theta. \tag{3.4}
 \end{aligned}$$

Now by Lemma 2.3, for each $\theta, 0 \leq \theta < 2\pi$ and any complex numbers $\alpha_1, \dots, \alpha_k, k \leq n-1$ with $|\alpha_i| \geq 1, i = 1, 2, \dots, k$ we have

$$\begin{aligned}
 & \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| \\
 & \leq \left| D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta}) \right| - mn(n-1) \cdots (n-k+1) (|\alpha_1 \cdots \alpha_k| - 1), \quad \text{for } |z| = 1.
 \end{aligned}$$

This implies

$$\begin{aligned}
 & \left\{ \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| + \frac{mn(n-1) \cdots (n-k+1) (|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\} \\
 & \leq \left\{ \left| D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta}) \right| - \frac{mn(n-1) \cdots (n-k+1) (|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\}. \tag{3.5}
 \end{aligned}$$

Take $A = |D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta})|$, $B = |D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta})|$,
 $C = \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2}$ in lemma 2.6, we get

$$B + C \leq A - C \leq A.$$

Hence for every real β , with the help of Lemma 2.6, we get

$$\begin{aligned} & \left| \left\{ |D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta})| - \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\} e^{i\beta} \right. \\ & \left. + \left\{ |D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta})| + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\} \right| \\ & \leq \left| |D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta})| e^{i\beta} + |D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta})| \right|. \end{aligned}$$

This implies for each $\gamma > 0$,

$$\int_0^{2\pi} \left| F(\theta) + e^{i\beta} G(\theta) \right|^\gamma d\theta \leq \int_0^{2\pi} \left| |D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta})| + e^{i\beta} |D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta})| \right|^\gamma d\theta, \tag{3.6}$$

where

$$F(\theta) = \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2}$$

and

$$G(\theta) = \left| D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta}) \right| - \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2}.$$

Integrating both sides of (3.6) with respect to β from 0 to 2π , we get with the help of (3.4), that for each $\gamma > 0$,

$$\begin{aligned} & \int_0^{2\pi} \int_0^{2\pi} \left| F(\theta) + e^{i\beta} G(\theta) \right|^\gamma d\theta d\beta \\ & \leq \int_0^{2\pi} \int_0^{2\pi} \left| |D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta})| + e^{i\beta} |D_{\alpha_1} \cdots D_{\alpha_k} Q(e^{i\theta})| \right|^\gamma d\theta d\beta \\ & \leq 2\pi n^\gamma (n-1)^\gamma \cdots (n-k+1)^\gamma (|\alpha_k| + 1)^\gamma \cdots (|\alpha_1| + 1)^\gamma \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta. \end{aligned} \tag{3.7}$$

Now for every real β and $t \geq 1$, we have

$$|t + e^{i\beta}| \geq |1 + e^{i\beta}|,$$

which implies for every $\gamma > 0$,

$$\int_0^{2\pi} |t + e^{i\beta}|^\gamma d\beta \geq \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta.$$

If $F(\theta) \neq 0$, we take $t = \left| \frac{G(\theta)}{F(\theta)} \right|$ and since $t \geq 1$ by (3.5)

$$\begin{aligned}
 & \int_0^{2\pi} |F(\theta) + e^{i\beta} G(\theta)|^\gamma d\beta \\
 &= |F(\theta)|^\gamma \int_0^{2\pi} \left| 1 + \frac{G(\theta)}{F(\theta)} e^{i\beta} \right|^\gamma d\beta \\
 &= |F(\theta)|^\gamma \int_0^{2\pi} \left| \frac{G(\theta)}{F(\theta)} + e^{i\beta} \right|^\gamma d\beta \\
 &= |F(\theta)|^\gamma \int_0^{2\pi} \left| \frac{G(\theta)}{F(\theta)} + e^{i\beta} \right|^\gamma d\beta \\
 &\geq |F(\theta)|^\gamma \int_0^{2\pi} \left| 1 + e^{i\beta} \right|^\gamma d\beta \\
 &= \left\{ \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\}^\gamma \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta.
 \end{aligned}$$

For $F(\theta) = 0$, this inequality is trivially true. Using this in (3.7), we conclude that for each $\gamma > 0, \beta$ real and any real or complex numbers $\alpha_1, \dots, \alpha_k, k \leq n-1$ with $|\alpha_i| \geq 1, i = 1, 2, \dots, k$,

$$\begin{aligned}
 & \int_0^{2\pi} |1 + e^{i\beta}|^\gamma d\beta \int_0^{2\pi} \left\{ \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2} \right\}^\gamma d\theta \\
 & \leq 2\pi n^\gamma (n-1)^\gamma \cdots (n-k+1)^\gamma (|\alpha_k| + 1)^\gamma \cdots (|\alpha_1| + 1)^\gamma \int_0^{2\pi} |P(e^{i\theta})|^\gamma d\theta.
 \end{aligned} \tag{3.8}$$

Now using the fact that for every real or complex number δ with $|\delta| \leq 1$,

$$\begin{aligned}
 & \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2} \delta \right| \\
 & \leq \left| D_{\alpha_1} \cdots D_{\alpha_k} P(e^{i\theta}) \right| + \frac{mn(n-1) \cdots (n-k+1)(|\alpha_1 \cdots \alpha_k| - 1)}{2},
 \end{aligned}$$

the desired result follows from (3.8).

References

- [1] Arestov, V. A., On Integral Inequalities for Trigonometric Polynomials and Their Derivatives, IZV., 18(1982), 1-17.
- [2] Aziz, A., A New Proof of Laguerre's Theorem About the Zeros of Polynomials, Bull. Austral. Math. Soc., 33 (1986), 133-138.
- [3] Aziz, A., Inequalities for the Polar Derivative of a Polynomial, J. Approx. Theory, 55 (1988), 183-193.
- [4] Aziz, A. and Dawood, Q. M., Inequalities for a Polynomial and its Derivative, J. Approx. Theory, 54 (1988), 306-313.

- [5] Aziz, A. and Rather, N. A., On an Inequality Concerning the Polar Derivative of a Polynomial, Proc. Indian Acad. Sci. Math. Sci., 117 (2003), 349-357.
- [6] Aziz, A. and Rather, N. A., New L^q Inequalities for Polynomials, Math. Ineq. and Appl., 1 (1998), 177-191.
- [7] De-Bruijn, N. G., Inequalities Concerning Polynomials in the Complex Domain, Nederl. Akad. Wetenschap. Proc., 50 (1947), 1265-1272.
- [8] Govil, N. K., Nyuydinkong, G. and Tameru, B., Some L^p Inequalities for the Polar Derivative of a Polynomial, J. Math. Anal. Appl., 254 (2001), 618-626.
- [9] Lax, P. D., Proof of a Conjecture of P. Erdős on the Derivative of a Polynomial, Bull. Amer. Math. Soc., 50 (1944), 509-513.
- [10] Marden, M., Geomertry of Polynomials, 2nd edition, Math. Surveys, No. 3, Amer. Math. Soc. (Providence, R.I.) (1966).
- [11] Milovanović, G. V., Mitrinović, D. S. and Rassias, Th. M., Topics in Polynomials, Extremal Problems, Inequalities, Zeros, World Scientific, Singapore, 1994.
- [12] Rahman, Q. I. and Schmeisser, G., L^p Inequalities for Polynomials, J. Approx. Theory, 53 (1988), 26-32.
- [13] Rather, N. A., L^p Inequalities for the Polar Derivative of a Polynomial, J. Ineq. Pure Appl. Math., 9 (2008), Issue 4, Art. 103, 10 pp.
- [14] Schaeffer, A. C., Inequalities of A. Markoff and S. Bernstein for polynomials and related functions, Bull. Amer. Math. Soc., 47 (1941), 565-579.
- [15] Zygmund, A., A Remark on Conjugate Series, Proc. London Math. Soc., 34(1932), 392-400.

Abdullah Mir

P. G. Department of Mathematics

Kashmir University

Hazratbal, Srinagar-190006

India

E-mail: mabdullah mir@yahoo.com

Sajad Amin Baba

Department of Mathematics

Govt. Hr. Sec. Institute

Kurhama Ganderbal-191201

India

E-mail: sajad2baba@yahoo.com