

## FRACTIONAL DIFFUSION EQUATIONS WITH INTERNAL DEGREES OF FREEDOM <sup>\*1)</sup>

Luis Vázquez

(*Departamento de Matemática Aplicada, Facultad de Informática, Universidad Complutense, E-28040 Madrid, Spain*)

(*Centro de Astrobiología CSIC-INTA, E-28850 Torrejón de Ardoz, Madrid, Spain*)

### Abstract

We present a generalization of the linear one-dimensional diffusion equation by combining the fractional derivatives and the internal degrees of freedom. The solutions are constructed from those of the scalar fractional diffusion equation. We analyze the interpolation between the standard diffusion and wave equations defined by the fractional derivatives. Our main result is that we can define a diffusion process depending on the internal degrees of freedom associated to the system.

*Key words:* Fractional derivative, Diffunors, Diffusion process, Generalized Dirac equation.

### 1. Introduction

It is well known the approach of Dirac to obtain his famous equation from the Klein-Gordon equation [1]. The free Dirac equation can be considered as the root square of the Klein-Gordon equation. In a more general context Morinaga and Nono [2] analyzed the integer  $s$ -root of the partial differential equations of the form

$$\sum_{|I|=s} a_I \frac{\partial^{|I|}}{\partial x^I} \phi = \phi \quad (1)$$

The  $s$ -root is the first order system

$$\sum_{i=1}^n \alpha_i \frac{\partial \Phi}{\partial x_i} = \Phi \quad (2)$$

being  $\alpha_1, \dots, \alpha_n$  matrices. From the physical point of view the  $\alpha_k$  describe internal degrees of freedom of the associated system.

The purpose of the paper is to generalize the above study to the case of fractional derivatives. In this context, we will consider the fractional diffusion equations with internal degrees of freedom obtained by generalization of the  $s$ -roots of the standard scalar linear diffusion equation. Thus, it is natural to consider the space and time fractional derivatives in a symmetric way through the framework of the standard Fourier transform

$$\frac{\partial^\alpha u(s)}{\partial s^\alpha} \longrightarrow (-i\kappa)^\alpha \hat{u}(\kappa) \quad (3)$$

being  $\kappa \in \mathfrak{R}$ .

In a future work we will consider the different following definitions of time and space fractional derivatives [3] which appear in other contexts:

---

\* Received March 20, 2001.

<sup>1)</sup>This work has been partially supported by the Project “The Sciences of Complexity” (ZiF, Bielefeld Universität), the Comisión Interministerial de Ciencia y Tecnología of Spain (grant PB98-0850) and the European Project COSIC of SENS (HPRN-CT-2000-00158).

- The time fractional derivative [4] of order  $\alpha > 0$  for a sufficiently well-behaved causal function  $u(t)$  is defined as follows

$$\frac{d^\alpha}{dt^\alpha}u(t) = \frac{1}{\Gamma(m - \alpha)} \int_0^t \frac{u^{(m)}(\tau)}{(t - \tau)^{\alpha+1-m}} d\tau \tag{4}$$

where  $m = 1, 2, \dots$ , and  $0 \leq m - 1 < \alpha < m$ . This definition requires the absolute integrability of the derivative of order  $m$ .

- The symmetric space fractional derivative [5] of order  $\alpha > 0$  of a sufficiently well-behaved function  $u(x), x \in \mathfrak{R}$ , is defined as the pseudo-differential operator characterized in its Fourier representation by

$$\frac{d^\alpha}{d|x|^\alpha}u(x) \longrightarrow -|\kappa|^\alpha \hat{u}(\kappa) \tag{5}$$

as before being  $\kappa \in \mathfrak{R}$ .

## 2. The Square Root of the Standard Linear Diffusion Equation

A possible definition of the root-square of the standard diffusion equation (SDE) in one space dimension,  $u_t - u_{xx} = 0$ , is the following:

$$\left(A \frac{\partial^{1/2}}{\partial t^{1/2}} + B \frac{\partial}{\partial x}\right)\psi(x, t) = 0 \tag{6}$$

where  $A$  and  $B$  are matrices satisfying the conditions:

$$A^2 = I \quad , \quad B^2 = -I \tag{7}$$

$$\{A, B\} \equiv AB + BA = 0 \tag{8}$$

being  $\psi(x, t)$  multidimensional with at least two scalar space-time components. Also, every scalar component satisfies the SDE. Such solutions can be interpreted as *probability distributions with internal structure associated to internal degrees of freedom of the system*. We could name them **diffunors** in analogy with the spinors in Quantum Mechanics.

We have two possible realizations of the above algebra in terms of real matrices  $2 \times 2$  associated to the Pauli matrices:

- 

$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad , \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tag{9}$$

- 

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad , \quad B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tag{10}$$

Other realizations involving complex bidimensional matrices are possible, but taking into account the reference to the diffusion equation we only consider the real representations.

The solutions of (6) are related to the SDE in a simple way. As an illustration, let us consider the representation (9), thus  $\psi(x, t) = \begin{pmatrix} \varphi(x, t) \\ \chi(x, t) \end{pmatrix}$

such that  $\chi(x, t) = \pm\varphi(x, t)$ . We have two general independent solutions of (6):

$$\varphi(x, t) \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \text{and} \quad \varphi(x, t) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \tag{11}$$

being  $\varphi(x, t)$  solution of SDE. The solutions (11) represent two possible probability distributions depending not only on the space and time coordinates, but also on the internal degrees of freedom. This effect could modelate the diffusion of particles with internal structure.

The equation (6) is not time reversible but it is invariant under space reflection as the underlying SDE. More precisely, in the representation given by (9) a possible representation of the **parity operator** is  $P = AP^0$ , such that  $P^0 : x \rightarrow -x$ .

### 3. The 1/3-root of the Standard Linear Diffusion Equation

As in the previous section, a possible definition of the 1/3-root of the SDE in one space dimension,  $u_t - u_{xx} = 0$ , is the following:

$$\left(M \frac{\partial^{1/3}}{\partial t^{1/3}} + N \frac{\partial^{2/3}}{\partial x^{2/3}}\right)\Psi(x, t) = 0 \tag{12}$$

where  $M$  and  $N$  are matrices satisfying the conditions:

$$M^3 = I, N^3 = -I \tag{13}$$

$$MMN + MNM + NMM = 0 \tag{14}$$

$$NNM + NMN + MNN = 0 \tag{15}$$

Being  $\Psi(x, t)$  multidimensional with at least three scalar space-time components. As before, we could name it **diffunors** with three components, in analogy with the spinors in Quantum Mechanics. A possible realization is in terms of the matrices  $3 \times 3$  associated to the Silvester algebra [6].

$$M = \begin{pmatrix} 0 & 0 & 1 \\ \omega^2 & 0 & 0 \\ 0 & \omega & 0 \end{pmatrix}, \quad N = \Omega \begin{pmatrix} 0 & 0 & 1 \\ \omega & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix} \tag{16}$$

where  $\omega$  is a cubic root of unity and  $\Omega$  a cubic root of the negative unity. As in the previous section, it can be made similar considerations about the solutions, but now we have three internal degrees of freedom.

### 4. General Time and Space-Fractional Diffusion Equations

Combining the results of the previous sections, we can consider the general fractional evolution equations:

$$\left(A \frac{\partial^\alpha}{\partial t^\alpha} + B \frac{\partial^\beta}{\partial x^\beta}\right)\Psi(x, t) = 0 \tag{17}$$

where the dynamics depends on the definition of the fractional derivatives, the orders  $\alpha$  and  $\beta$  and on the structure of the matrices  $A$  and  $B$ . We can consider the following general considerations:

- If  $A$  and  $B$  are the identity, the equation (17) defines a sort of interpolation between the standard diffusion equation,  $u_t - u_{xx} = 0$ , and the standard wave equation,  $u_{tt} - u_{xx} = 0$ , while  $0 < \alpha \leq 2$  and  $0 < \beta \leq 2$ . The solutions are well studied in [7]-[9] when the definition of fractional derivatives (4)-(5) are considered.

When  $\beta = 2$ , the absolute moments of even order are the following

$$\langle x^{2n} \rangle = \frac{\Gamma(2n + 1)}{\Gamma(\alpha n + 1)} t^{\alpha n} \tag{18}$$

being  $n = 0, 1, 2, \dots$

- If we consider the algebraic structure defined by (8) with  $0 < \alpha \leq 1$  and  $0 < \beta \leq 1$ , we define other alternative to the above interpolation through the Dirac ( $\alpha = 1, \beta = 1$ ) equation which includes the effects associated to the internal degrees of freedom. In this case, the solution of the equation is related to the solutions of the above fractional diffusion-wave equations when  $A = B = I$ .
- With the definition (3) of fractional derivatives, the absolute moments of even order for each one of the two degrees of freedom of (6) are the following

$$\langle x^{2n} \rangle = \frac{\Gamma(2n+1)}{\Gamma(n+1)} t^n \quad (19)$$

with  $n = 0, 1, 2, \dots$ , since the associated functions satisfy the standard diffusion equation.

### References

- [1] S.S. Schweber, *An Introduction to Relativistic Quantum Field Theory*, Harper Int. Ed., New York, 1966.
- [2] K. Morinaga and T. Nono, *J. Sci. Hiroshima Univ.(A)*, **16** (1952), 13-41.
- [3] B. Ross (Ed.), *Fractional Calculus and its Applications*, Lecture Notes in Mathematics **457**, Springer-Verlag, 1975.
- [4] M. Caputo and F. Mainardi, Linear models of dissipation in anelastic solids, *Riv. Nuovo Cimento (Ser.II)*, **1** (1971), 161-198.
- [5] A.I. Saichev and G.M. Zaslavsky, Fractional kinetic equations: solutions and applications, *Chaos* **7** (1997), 753-764.
- [6] N. Heerema, *Duke Math. J.*, **21** (1954), 423-443.
- [7] F. Mainardi, *Fractals and Fractional Calculus in Continuum Mechanics*, Eds. A. Carpinteri and F. Mainardi, CISM Courses and Lectures **378**, Springer-Verlag, Wie, 291-348 (1997).
- [8] F. Mainardi and M. Tomirotti, Seismic pulse propagation with constant Q and stable probability distributions, *Annali di Geofisica*, **40** (1997), 1311-1328.
- [9] R. Gorenflo and F. Mainardi, Fractional calculus and stable probability distributions, *Arch. Mechanics*, **50** (1998), 377-388.