

First-Principles Calculations of Shocked Fluid Helium in Partially Ionized Region

Cong Wang¹, Xian-Tu He^{1,2} and Ping Zhang^{1,2,*}

¹ LCP, Institute of Applied Physics and Computational Mathematics, P.O. Box 8009, Beijing 100088, China.

² Center for Applied Physics and Technology, Peking University, Beijing 100871, China.

Received 29 April 2011; Accepted (in revised version) 12 December 2011

Communicated by Xingao Gong

Available online 17 April 2012

Abstract. Quantum molecular dynamic simulations have been employed to study the equation of state (EOS) of fluid helium under shock compressions. The principal Hugoniot is determined from EOS, where corrections from atomic ionization are added onto the calculated data. Our simulation results indicate that principal Hugoniot shows good agreement with gas gun and laser driven experiments, and maximum compression ratio of 5.16 is reached at 106 GPa.

AMS subject classifications: 81Q05, 82D10, 85A04

Key words: Equation of state, warm dense matter, ab initio simulations.

1 Introduction

High-pressure introduced response of materials, which requires accurate understandings of the thermophysical properties into new and complex region, has gained much scientific interest recently [1]. The relative high temperature and high density are usually referred as the so-called "warm dense matter" (WDM)—a strongly correlated state, where simultaneous dissociations, ionizations, and degenerations make modelling of the dynamical, electrical, and optical properties of WDM extremely challenging [2]. WDM, which provides an active research platform by combining the traditional plasma physics

*Corresponding author. *Email addresses:* wang_cong@iapcm.ac.cn (C. Wang), xthe@iapcm.ac.cn (X.-T. He), zhang_ping@iapcm.ac.cn (P. Zhang)

and condensed matter physics, usually appears in shock or laser heated targets [3], inertial confinement fusion [4], and giant planetary interiors [5].

Next to hydrogen, helium is the most abundant element in the universe, and physical properties of warm dense helium, especially the EOS, are critical for astrophysics [6, 7]. For instance, the structure and evolution of stars, White Dwarfs, and Giant Planets [8–11], and therefore the understanding of their formation, depends sensitively on the EOS of hydrogen and helium at several megabar regime. For all planetary models, accurate EOS data are essential in solving the hydrostatic equation. As a consequence, a series of experimental measurements and theoretical approaches have been applied to investigate the EOS of helium. Liquid helium was firstly single shocked to 15.6 GPa using two stage light gas gun by Nellis *et al.*, then double shocked to 56 GPa, and the calculated temperature are 12000 and 22000 K respectively [12]. Maximum compression ratio ($\eta_{max} \approx 6$) was achieved by laser driven shock experiments with the crossover pressure around 100 GPa [13], However, soon after that, Knudson *et al.* have modified η_{max} to be 5.1 [14]. Theoretically, since the ionization equilibrium is not interfered with the dissociation equilibrium between molecules and atoms, helium, which is characterized by monoatomic molecule and close shell electronic structure, is particularly suitable for the investigation of the high pressure behavior under extreme conditions. Free energy based chemical models by Ross *et al.* [15], Chen *et al.* [16], and Kowalski *et al.* [17] have been used to investigate the principal Hugoniot of liquid helium, and the results are accordant with gas gun experiments. However, considerable controversies have been raised at megabar pressure regime, especially since the data was probed by laser shock wave experiments. Interatomic potential method predict $\eta_{max}=4$ for shock compressed helium [15], whereas, the EOS used by the astrophysical community from Saumon *et al.* (SCVH) [8], path-integral Monte Carlo (PIMC) [18], and activity expansion (ACTEX) [19] calculations provide an increase in compressibility at the beginning of ionization. For the initial density of $\rho_0=0.1233 \text{ g/cm}^3$, SCVH and ACTEX simulation results indicate the maximum compression ratio lies around 6 at 300 GPa and 100 GPa respectively [8, 19], while PIMC calculations suggest $\eta_{max}=5.3$ near 360 GPa [18].

On the other hand, quantum molecular dynamic (QMD) simulations, where quantum effects are considered by the combinations of classical molecular dynamics for the ions and density functional theory (DFT) for electrons, have already been proved to be successful in describing thermophysical properties of materials at complex conditions [20, 21]. However, the DFT based molecular dynamic simulations (with or without accounting for excited electrons) do not provide reasonable results at the ionization region, mainly because the atomic ionization is not well defined in the framework of DFT. Considering these facts mentioned above, thus, in the present work, we applied the corrected QMD simulations to study shock compressed helium, and the EOS, which is compared with experimental measurements and different theoretical models, are determined for a wide range of densities and temperatures. The calculated compression ratio is substantially increased according to the ionization of atoms in the warm dense fluid.

2 Computational method

The Vienna Ab-initio Simulation Package (VASP) [22, 23], which was developed at the Technical University of Vienna, has been employed to perform simulations for helium. The elements of our calculations consist of a series of volume-fixed supercells including N atoms, which are repeated periodically throughout the space. By involving Born-Oppenheimer approximation, electrons are quantum mechanically treated through plane-wave, finite-temperature (FT) DFT [24], where the electronic states are populated according to Fermi-Dirac distributions at temperature T_e . The exchange-correlation functional is determined by generalized gradient approximation (GGA) with the parametrization of Perdew-Wang 91 [25]. The ion-electron interactions are represented by a projector augmented wave (PAW) pseudopotential [26]. Isokinetic ensemble (NVT) is adopted in present simulations, where the ionic temperature T_i is controlled by Noé thermostat [27], and the system is kept in local equilibrium by setting the electron (T_e) and ion (T_i) temperatures to be equal.

The plane-wave cutoff energy is selected to be 700.0 eV so that the pressure is converged within 3% accuracy. Γ point is used to sample the Brillouin zone in molecular dynamics simulations, because EOS can only be modified within 5% for the selection of higher number of \mathbf{k} points. 64 helium atoms are included in the cubic supercell. The densities selected in our simulations range from 0.1233 to 0.8 g/cm³ and temperatures between 4 and 50000 K, which highlight the regime of the principal Hugoniot. All the dynamic simulations are lasted for 6000 steps, and the time steps for the integrations of atomic motion are selected according to different densities (temperatures) [28]. Then, the subsequent 1000 steps of simulation are used to calculate EOS as running averages.

3 Results and discussion

Different corrections to QMD simulations have already been used to model the thermo-physical properties of WDM, such as the zero point vibrational energy modification for hydrogen and deuterium [29]. However, quantitative descriptions of the ionization of atoms in the frame of DFT is still lacking, except for the Drude model for aluminium introduced by Mazevet *et al.* [30], but the simple metallic model is not suitable for studying warm dense helium. Here, we adopt similar approximations as described in [31], where the effect of atomic ionization are considered, to study the EOS of fluid helium. In this work, the EOS are expressed as follows [31]:

$$E = E_{QMD} + N\beta E_{ion}, \quad (3.1)$$

$$P = P_{QMD} + (1 + \beta) \frac{\rho k_B T}{m_{He}}, \quad (3.2)$$

where E_{QMD} and P_{QMD} are data obtained from regular QMD calculations, and N is the total number of atoms considered in supercells. m_{He} and k_B represent the mass of helium

atom and Boltzmann constant. The density and temperature are denoted by ρ and T , respectively. β stands for the ionization degree, and E_{ion} is the ionization energy. Van der Waals interactions are treated as small as negligible in the present work.

In the present work, only the first ionization level is considered, and the ionization degree of helium can be evaluated through Saha equation:

$$\frac{\beta^2}{1-\beta} = \frac{2\Omega}{\lambda^3} \exp\left(-\frac{E_{ion}}{k_B T}\right), \quad (3.3)$$

$$\lambda = \sqrt{\frac{h^2}{2\pi m_e k_B T}}, \quad (3.4)$$

where, m_e and Ω present the mass of electron and volume of the supercell. The ionization energy E_{ion} is determined by $E_{ion} = E^{He^+}(\rho, T) - E^{He}(\rho, T)$, where $E^{He^+}(\rho, T)$ and $E^{He}(\rho, T)$ are total energy of He^+ and He at the relative density and temperature, respectively. These energies are calculated from FT-DFT.

Based on the approximations mentioned above, the calculated EOS was examined theoretically along the principal Hugoniot, the locus of states that satisfy the Rankine-Hugoniot (RH) equations, which are derived from conservation of mass, momentum, and energy across the front of shock waves. The RH equations describe the locus of states in (E, P, V) -space satisfying the following relation:

$$E_1 - E_0 = \frac{1}{2}(P_1 + P_0)(V - V_0), \quad (3.5)$$

$$P_1 - P_0 = \rho_0 u_s u_p, \quad (3.6)$$

$$V_1 = V_0(1 - u_p / u_s), \quad (3.7)$$

where E, P, V present internal energy, pressure, volume, and subscripts 0 and 1 present the initial and shocked state, respectively. In Eqs. (3.6) and (3.7), u_s is the velocity of the shock wave and u_p corresponds to the mass velocity of the material behind the shock front. In our present simulations, the initial density for helium is $\rho_0 = 0.1233 \text{ g/cm}^3$ and the liquid specimen is controlled at a temperature of 4 K, where the internal energy is $E_0 = -0.02 \text{ eV/atom}$. The initial pressure P_0 can be treated approximately as zero compared to the high pressure of shocked states along the Hugoniot. The Hugoniot points are obtained as follows: (i) smooth functions are used to fit the internal energy and pressure in terms of temperature at sampled density; (ii) then, Hugoniot points are derived from Eq. (3.5).

The principal Hugoniot curve for helium is shown in Fig. 1, where previous experimental measurements and theoretical predictions are also plotted for comparison. For pressures below 30 GPa, our results show good agreements with the data detected by gas gun experiments [12]. At higher pressures, the principal Hugoniot curve shows a maximum compression ratio $\eta_{max} \approx 6$ with the crossover pressure around 100 GPa, as have been reported by laser driven experiments [13]. Soon after that, the use of quartz




Figure 1: (Color online) Principal Hugoniot up to 130 GPa is shown for the present work (red line), where previous theoretical predictions and experimental measurements are also shown for comparison. Experiments: Two stage light gas gun data [12] is denoted by open circle; Laser driven experimental results [13] are labeled by open square. Theories: (i) Classical MC simulation results [32] are plotted as green solid line; (ii) ideal plasma's Hugoniot curve [18] is shown as green dotted line; (iii) SCVH method [8] is labeled as green dashed line; (iv) ACTEX results [19] are blue solid line; (v) Previous DFT simulation results [18] (with and without accounting for electron excitation) are shown as black dashed and solid line, respectively; (vi) PIMC results [18] are labeled as black dotted line. Inset is the plot of shock wave velocity verse mass velocity.

as a shock wave standard by Knudson *et al.* has improve the shock data, and η_{max} has been reduced to be 5.1 [14]. Our corrected QMD simulation results indicate that, the atomic ionization dominates the characteristic of the EOS at $P > 30$, and the principal Hugoniot shows soften behavior with the increase of pressure, then reach its maximum ($\eta_{max}=5.16$) with the corresponding pressure of 106 GPa. Then, stiff behavior has been found at $P > 110$ GPa along the Hugoniot. As has been shown in Fig. 2, with the increase




Figure 2: Ionization energy and ionization degree are plotted as functions of pressure along the principal Hugoniot in the upper and lower panels, respectively.




Figure 3: (Color online) Comparison between our calculated shock temperature and previous theoretical predictions. Interatomic potential results are shown as open square [12] and solid line [15]; DFT [18] and ACTEX [19] results are plotted as dashed and dotted line, respectively.

of pressure along the Hugoniot, ionization energy decreases, and continuous increase of atomic ionization has become considerable in determining the EOS of fluid helium. Maximum compression ratio is achieved with the ionization degree of 40%. The wide-range behavior of our simulated principal Hugoniot for helium shows excellent agreement with experimental ones.

Concerning other theoretical models, classical MC results show stiff behavior along the Hugoniot, and η_{max} lies around 4 at 1000 GPa [32]. Direct DFT simulation results show stiff behavior up to 110 GPa, while, by combining PIMC simulations, η_{max} was reported to be 5.3 at 360 GPa [18], but the predicted pressure is still too high compared with laser driven experiments. ACTEX [19] and SCVH [8] calculations predict η_{max} to be 6 near 100 GPa and 300 GPa respectively, but the results do not agree with those of Knudson *et al.* [14].

Temperature, which is focused as one of the most important parameters in experiments, is difficult to be measured because of the uncertainty in determining the optical-intensity loss for ultraviolet part of the spectrum in adiabatic or isentropic shock compressions, especially for the temperature exceeding several electron-volt [33]. QMD simulations provide powerful tools to predict shock temperature. The calculated Hugoniot temperature has been shown in Fig. 3 as a function of pressure along the Hugoniot. Previous theoretical predictions, such as interatomic potential [12,15], DFT-MD (excited electron) [18], and ACTEX [19] are also provided for comparison. Disagreements have been found to begin at around 30 GPa with the relative temperature of 20000 K, which highlights the starting point of ionization on the Hugoniot, and the predicted temperatures by those models are higher at a given pressure compared with our calculations.

4 Conclusion

In summary, We have performed DFT based QMD simulations to study the thermophysical properties of helium under extreme conditions. The Hugoniot EOS has been evaluated through QMD calculations and corrected by taking into account the atomic ionization described by Saha equation, where only first ionization is considered. The corrected Hugoniot has been proved accord well with the experimental data in a wide range of shock conditions, which thus indicates the importance of atomic ionization. Maximum compression ratio of 5.16 reveals at round 106 GPa along the principal Hugoniot, and the softened characteristic of the Hugoniot has been demonstrated by the contributions from

Acknowledgments

This work was supported by NSFC under Grant No. 11005012, by the National Basic Security Research Program of China, and by the National High-Tech ICF Committee of China.

References

- [1] R. Ernstorfer, M. Harb, C. T. Hebeisen, G. Sciaini, T. Dartigalongue, and R. J. D. Miller, The formation of warm dense matter: Experimental evidence for electronic bond hardening in gold, *Science*, 323 (2009), 1033-1037.
- [2] W. J. Nellis, Dynamic compression of materials: metallization of fluid hydrogen at high pressures, *Rep. Prog. Phys.*, 69 (2006), 1479.
- [3] D. G. Hicks, T. R. Boehly, P. M. Celliers, J. H. Eggert, S. J. Moon, D. D. Meyerhofer, and G. W. Collins, Laser-driven single shock compression of fluid deuterium from 45 to 220 GPa, *Phys. Rev. B*, 79 (2009), 014112.
- [4] F. Philippe, A. Casner, T. Caillaud, O. Landoas, M. C. Monteil, S. Liberatore, H. S. Park, P. Amendt, H. Robey, C. Sorce, C. K. Li, F. Seguin, M. Rosenberg, R. Petrasso, V. Glebov, and C. Stoeckl, Experimental demonstration of X-ray drive enhancement with rugby-shaped hohlraums, *Phys. Rev. Lett.*, 104 (2010), 035004.
- [5] W. Lorenzen, B. Holst, and R. Redmer, Demixing of hydrogen and helium at megabar pressures, *Phys. Rev. Lett.*, 102 (2009), 115701.
- [6] D. J. Stevenson and E. E. Salpeter, The phase diagram and transport properties for hydrogen-helium fluid planets, *Astrophys. J. Suppl.*, 35 (1977), 221-237.
- [7] D. J. Stevenson and E. E. Salpeter, The dynamics and helium distribution in hydrogen-helium fluid planets, *Astrophys. J. Suppl.*, 35 (1977), 239-261.
- [8] D. Saumon, G. Chabrier, and H. M. Van Horn, An equation of state for low-mass stars and giant planets, *Astrophys. J. Suppl. Ser.*, 99 (1995), 713-41.
- [9] V. Ya. Ternovoi, S. V. Kvitov, A. A. Pyalling, A. S. Filimonov and V. E. Fortov, Experimental determination of the conditions for the transition of Jupiters atmosphere to the conducting state, *JETP Lett.*, 79 (2004), 6-9.
- [10] J. Vorberger, I. Tamblyn, B. Militzer, and S. A. Bonev, Hydrogen-helium mixtures in the interiors of giant planets, *Phys. Rev. B*, 75 (2007), 024206.

- [11] M. A. C. Perryman, Extra-solar planets, *Rep. Prog. Phys.*, 63 (2000), 1209-1272.
- [12] W. J. Nellis, N. C. Holmes, A. C. Mitchell, R. J. Trainor, G. K. Governo, M. Ross, and D. A. Young, Shock Compression of liquid helium to 56 GPa (560 kbar), *Phys. Rev. Lett.*, 53 (1984), 1248.
- [13] J. Eggert, S. Brygoo, P. Loubeyre, R. S. McWilliams, P. M. Celliers, D. G. Hicks, T. R. Boehly, R. Jeanloz, and G.W. Collins, Hugoniot data for helium in the ionization regime, *Phys. Rev. Lett.*, 100 (2008), 124503.
- [14] M. D. Knudson and M. P. Desjarlais, Shock compression of quartz to 1.6 TPa: Redefining a pressure standard, *Phys. Rev. Lett.*, 103 (2009), 225501.
- [15] M. Ross and D. A. Young, Helium at high density, *Phys. Lett. A*, 118 (1986), 463-466.
- [16] Q. F. Chen, Y. Zhang, L. C. Cai, Y. J. Gu, and F. Q. Jing, Self-consistent variational calculation of the dense fluid helium in the region of partial ionization, *Phys. Plasma.*, 14 (2007), 012703.
- [17] P. M. Kowalski, S. Mazevet, D. Saumon, and M. Challacombe, Equation of state and optical properties of warm dense helium, *Phys. Rev. B*, 76 (2007), 075112.
- [18] B. Militzer, First principles calculations of shock compressed fluid helium, *Phys. Rev. Lett.*, 97 (2006), 175501.
- [19] M. Ross, F. Rogers, N. Winter, and G. Collins, Activity expansion calculation of shock-compressed helium: The liquid Hugoniot, *Phys. Rev. B*, 76 (2007), 020502(R).
- [20] C. Wang and P. Zhang, Ab initio study of shock compressed oxygen, *J. Chem. Phys.*, 132 (2010), 154307.
- [21] C. Wang and P. Zhang, The equation of state and nonmetal-metal transition of benzene under shock compression, *J. Appl. Phys.*, 107 (2010), 083502.
- [22] G. Kresse and J. Hafner, Ab initio molecular dynamics for liquid metals, *Phys. Rev. B*, 47 (1993), R558.
- [23] G. Kresse and J. Furthmüller, Efficient iterative schemes for ab initio total-energy calculations using a plane-wave basis set, *Phys. Rev. B*, 54 (1996), 11169.
- [24] T. Lenosky, S. Bickham, J. Kress, and L. Collins, Density-functional calculation of the Hugoniot of shocked liquid deuterium, *Phys. Rev. B*, 61 (2000), 1.
- [25] J. P. Perdew, *Electronic Structure of Solids*, Akademie Verlag, Berlin (1991).
- [26] P. E. Blöchl, Projector augmented-wave method, *Phys. Rev. B*, 50 (1994), 17953.
- [27] S. Nosé, A unified formulation of the constant temperature molecular dynamics methods, *J. Chem. Phys.*, 81 (1984), 511.
- [28] The time steps have been taken as $\Delta t = a/20\sqrt{k_B T/m_{He}}$, where $a = (3/4\pi n_i)^{1/3}$ is the ionic sphere radius (n_i is the ionic number density), $k_B T$ presents the kinetic energy, and m_{He} is the ionic mass.
- [29] B. Holst, R. Redmer, and M. P. Desjarlais, Thermophysical properties of warm dense hydrogen using quantum molecular dynamics simulations, *Phys. Rev. B*, 77 (2008), 184201.
- [30] S. Mazevet, M. P. Desjarlais, L. A. Collins, J. D. Kress, and N. H. Magee, Simulations of the optical properties of warm dense aluminum, *Phys. Rev. E*, 71 (2005), 016409.
- [31] C. Wang, X. T. He, and P. Zhang, Hugoniot of shocked liquid deuterium up to 300 GPa: Quantum molecular dynamic simulations, *J. Appl. Phys.*, 108 (2010), 044909.
- [32] R. A. Aziz, A. R. Janzen, and M. R. Moldover, Ab initio calculations for helium: A standard for transport property measurements, *Phys. Rev. Lett.*, 74 (1995), 1586.
- [33] Q. F. Chen, private communication, (2010).